

TO CAUSE A HAZARD IN THE USE, STORAGE OR DISPOSAL OF THIS SUBSTANCE IS AN OFFENCE.

PRECAUTIONS:

1. Handle with care. Harmful when swallowed. Avoid contact. Dangerous when inhaled. **Highly corrosive.**
2. Store away from sun and damp conditions.
3. Store in the original closed container in a cool, dry and well-ventilated place **UNDER LOCK AND KEY.**
4. **KEEP APART FROM FOOD AND FOODSTUFFS, SEEDS AND FERTILIZERS.**
5. **KEEP OUT OF REACH OF CHILDREN, UNAUTHORIZED PERSONS AND ANIMALS.**
6. **WEAR PROTECTIVE CLOTHING E.G. OVERALL, RUBBER GLOVES, NOSE COVERS AND GUMBOOTS WHEN HANDLING, PREPARING AND APPLYING THE SPRAY MIXTURE.**
7. Avoid prolonged or repeated exposure.
8. **DO NOT** carry **ZeroToll** in a vehicle which is used to carry people.
9. Toxic to aquatic organisms.
10. Wash contaminated clothing after use.
11. Wash with soap and water after use or after accidental skin contact.
12. Avoid eye and skin contact and inhalation of spray mist.
13. Do not eat, drink or smoke while mixing, applying or before washing hands and face or change of clothing.
14. **DO NOT** contaminate streams, rivers or waterways with the chemical or used containers.
15. Clean application equipment after use. Do not dispose of water where it can contaminate other crops, grazing, rivers or dams, feedstuffs, drinking water and eating utensils.

CONTAINER DISPOSAL: Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances.

DEALING WITH SPILT PRODUCT: In case of spill, flood area with large quantities of water. Do not store in a manner where cross-contamination with other pesticides or fertilizers could occur.

SYMPTOMS OF POISONING: DANGER. CORROSIVE. Causes skin burns. Causes irreversible eye damage. Harmful if swallowed. Harmful if inhaled. Avoid breathing vapor or spray mist. Do not get in eyes, on skin, or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

FIRST AID INSTRUCTIONS:

If on skin or clothing: Take off contaminated clothing and rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control center or doctor for treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call a poisons information center or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

USE ONLY ACCORDING TO LABEL INSTRUCTIONS. CAREFULLY READ THE ENTIRE LABEL BEFORE USE INCLUDING THE ACCOMPANYING LEAFLET.

DISCLAIMER: Although this remedy has been extensively tested under a large variety of conditions, the supplier does not warrant that it will be efficacious under all conditions, because the action and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions, quality of diluted water, compatibility with other substances not indicated on the label and the occurrence of resistance of the weeds to the remedy concerned, as well as by the method, time and accuracy of application. The supplier furthermore does not accept responsibility for performance of the remedy concerned due to failure of the user to follow label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier.

ZeroToll



HARMFUL BY SKIN CONTACT

Composition (weight by volume):

- Hydrogen Peroxide: 27%
- Peroxyacetic Acid: 2.0%

Chemical Groups:

- Hydrogen Peroxide: Peroxides
- Peroxyacetic Acid: Organic peroxide

Net Content:
20Lt

Personal protection equipment is mandatory for ZeroToll use, due to its high corrosivity

ZeroToll is a highly active, fungicide and bactericide against diseases on a wide range of crops

Manufactured for:



SINERIA Holland BV
Randwycksingel 20-A015
6229 EE Maastricht,
The Netherlands
www.sineria.com

Shelf Life: Two years from manufacturing date when stored in unopened, original container in a cool, dry and well-ventilated place

Manufacture Date:	
Expiration Date:	
Batch Number:	

HIGHLY CORROSIVE CONTENT
READ THE LABEL BEFORE USING THE PRODUCT
KEEP OUT OF THE REACH OF CHILDREN



18UAJH

GENERAL INFORMATION:

DESCRIPTION: ZeroToll is a highly active, fungicide and bactericide against diseases on a wide range of crops.
DIRECTIONS FOR USE: Do not apply this product through any irrigation system unless the chemigation instructions on this label are followed. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Application rates and instructions:

Crop	Disease	Application Method	Rate	Directions
Cucurbit: Cucumber, Melons, Pumpkin, Squash	Root and Fruit Rot (<i>Phytophthora</i> spp.), Downy Mildew, Powdery Mildew, Angular Leaf Spot	Foliar Spray	5.0-6.0 Lt/ha	As a foliar spray, apply product as a dilute spray (not to exceed 2.5% v/v (1:40) concentration in water). Ensure complete coverage of foliage. Repeat applications on 7–14 day interval.
		Irrigation (Overhead or Drip)		When applying through irrigation, make first application immediately after planting and repeat applications on 14–21 day interval.
Fruiting Vegetables: Eggplant, Peppers, Tomato	Late Blight and Root Rot (<i>Phytophthora infestans</i> and <i>Phytophthora</i> spp.), Powdery Mildew, Bacterial Spot and Speck, Bacterial Wilt	Foliar Spray	5.0-6.0 Lt/ha	As a foliar spray, apply product as a dilute spray (not to exceed 2.5% v/v (1:40) concentration in water). Ensure complete coverage of foliage. Repeat applications on 7–14 day interval.
		Irrigation (Overhead or Drip)		When applying through irrigation, make first application immediately after planting and repeat applications on 14-21 day interval.
Legumes: Chick Peas, Dry Beans, Lima Beans, Peas, Snap Beans	Bacterial Blight, Powdery Mildew, <i>Pythium</i> spp., Downy Mildew	Foliar Spray	4.0-6.0 Lt/ha	Apply product as a dilute spray (not to exceed 2.5% v/v (1:40) concentration in water). Ensure complete coverage of foliage. Repeat applications on 7–14 day interval.
Potato Pre-Harvest	Late Blight (<i>Phytophthora infestans</i>) Pink Rot (<i>Phytophthora erythroseptica</i>) Pythium Leak (<i>Pythium</i> spp.)	Foliar Spray	5.0-6.0 Lt/ha	As a foliar spray, apply once every 7–14 days depending on disease conditions. Product can be rotated with other fungicides labeled for Late Blight.
		In-furrow	5.0-8.0 Lt/ha	As in-furrow application, apply in a band directly over the seed pieces prior to row closure using 50–100Lt of water per ha. Additional in-season foliar applications may be necessary.

ORNAMENTALS:

Disease	Application Method	Dilution Rate	Application Rate	Directions
Downy Mildew	Foliar Spray	1:100-1:300	1.5-5.0 Lt per 500 Lt of water	Apply at first sign of disease and repeat as needed at 7–14 day interval. Ensure thorough wetness of all foliage. Use the higher rate when disease pressure is severe.
	Soil Drench	1:500-1:1,000	15.0-30.0 Lt per 500 Lt of water	As a drench, apply 1.5-3.0 Lt of mixed solution per square meter. Repeat as necessary.
Bacterial Blight, Bacterial Leaf Spot, <i>Phytophthora</i> spp. and <i>Pythium</i> spp.	Foliar Spray	1:100-1:300	1.5-5.0 Lt per 500 Lt of water	Apply at first sign of disease and repeat as needed at 7–14 day interval. Ensure thorough wetness of all foliage. Use the higher rate when disease pressure is severe.
	Soil Drench	1:500-1:1,000	15.0-30.0 Lt per 500 Lt of water	As a drench, apply 1.5-3.0 Lt of mixed solution per square meter. Repeat as necessary.
	Transplant Dip	1:500	3.0 Lt per 50 Lt of water	Dip the transplants for at least 30 seconds immediately before transplanting. Ensure root mass is thoroughly wet.

COMPATIBILITY: The supplier makes no recommendation in regard to the compatibility of this product with any other products. No warranty is provided by the supplier for mixtures of this product with any other products. When mixing this product with other products, the buyer / user takes full responsibility and risk.

WARRANTY: Seller's guarantee is limited to the terms set out on the label and subject there to, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the products on that condition.

IMPORTANT NOTE: This product must be used strictly as directed and in accordance with the instructions appearing on the label. The supplier accepts no liability other than as indicated in the Supplier's 'terms and conditions of sale' and the warranty clauses indicated on the product's label.

Distributor:



Cropserve (Pvt) Ltd
23 Anthony Ave, Msasa,
Harare, Zimbabwe
Phone: +263 242 307 628/9
Email: info@cropserve.co.zw
www.cropserve.co.zw